

# ON THE KAUFFMAN-HARARY CONJECTURE FOR ALEXANDER QUANDLE COLORINGS

SOICHIRO ASAMI

ABSTRACT. The Kauffman-Harary conjecture is a conjecture for Fox's colorings of alternating knots with prime determinants. We consider a conjecture for Alexander quandle colorings by referring to the Kauffman-Harary conjecture. We prove that this new conjecture is true for twist knots.

## 1. INTRODUCTION.

We recall the following conjecture called the *Kauffman-Harary conjecture* [3].

**Conjecture 1.1.** *Let  $D$  be a reduced alternating knot diagram with a prime determinant  $p$ . Then every non-trivial Fox's  $p$ -coloring of  $D$  assigns different colors to different arcs of  $D$ .*

In [1], Asaeda, Przytycki and Sikora generalize the conjecture by stating it in terms of homology of the double cover of the 3-sphere  $S^3$  branched along a link, and prove that the generalized conjecture is true for Montesinos links.

Fox's  $p$ -coloring is coincident with the coloring by the dihedral quandle  $R_p$  of order  $p$  (cf. [2]). We consider the following conjecture associated with the Alexander quandle which we can regard as a generalization of  $R_p$ .

**Conjecture 1.2.** *Let  $K$  be an alternating oriented knot,  $D$  be a reduced alternating diagram of  $K$ , and  $\Delta_K(t)$  be the Alexander polynomial of  $K$ . If the ring  $\mathbb{Z}[t, t^{-1}]/(\Delta_K(t))$  is an integral domain, then every non-trivial coloring of  $D$  by the Alexander quandle  $\mathbb{Z}[t, t^{-1}]/(\Delta_K(t))$  assigns different colors to different arcs of  $D$ .*

If the Kauffman-Harary conjecture is true then Conjecture 1.2, also, is true for any alternating knots with prime determinants (see Section 2). But Conjecture 1.2 is not included in the Kauffman-Harary conjecture because there is a knot with a non-prime determinant whose Alexander polynomial is a prime element in the Laurent polynomial ring  $\mathbb{Z}[t, t^{-1}]$ .

This paper is organized as follows. In Section 2, we review the definition of quandles and colorings by quandles of knot diagrams. In Section 3, we study colorings by Alexander quandles for the diagram of twist knots. Finally, we prove that Conjecture 1.2 is true for twist knots (Theorem 3.3).

## 2. QUANDLES AND COLORINGS.

In this section, we review the definition of quandles and colorings by quandles.

**Definition 2.1** ([4], [5]). A quandle,  $X$ , is a set with a binary operation  $*$  :  $X \times X \rightarrow X$  satisfying the following conditions:

---

2000 *Mathematics Subject Classification.* 57Q25, 57M42.

- (Q1) For any  $x \in X$ ,  $x * x = x$ .  
(Q2) For any  $x, y \in X$ , there is a unique element  $z \in X$  such that  $x = z * y$ .  
(Q3) For any  $x, y, z \in X$ ,  $(x * y) * z = (x * z) * (y * z)$ .

The condition (Q2) is equivalent to the following condition:

- (Q2') For any  $x, y \in X$ , there is a binary operation  $*^{-1} : X \times X \rightarrow X$  such that  $(x * y) *^{-1} y = x = (x *^{-1} y) * y$ .

We list some typical examples of quandles.

- Example 2.2.** (1) Let  $m$  be a positive integer, and for elements  $i, j \in \mathbb{Z}_m$ , define  $i * j = 2j - i \pmod{m}$ . Then the operation  $*$  defines a quandle structure called the *dihedral quandle*  $R_m$  of order  $m$ . The operation  $*^{-1}$  is identical with the operation  $*$ .  
(2) Let  $\Lambda = \mathbb{Z}[t, t^{-1}]$  be the Laurent polynomial ring over  $\mathbb{Z}$ ,  $J \subset \Lambda$  be an ideal of  $\Lambda$ . Then the quotient ring  $\Lambda/J$  with the binary operation defined by  $x * y = tx + (1 - t)y$  for any  $x, y \in \Lambda/J$  is a quandle called an *Alexander quandle*. The operation  $*^{-1}$  is given by  $x *^{-1} y = t^{-1}x + (1 - t^{-1})y$ . We remark that the dihedral quandle  $R_m$  is isomorphic to  $\Lambda/(m, t + 1)$ .

**Definition 2.3.** Let  $D$  be a diagram of an oriented knot  $K$ , and  $\Sigma$  the set of arcs of  $D$ . Given a quandle  $X$ , an  $X$ -coloring for  $D$  is a map  $C : \Sigma \rightarrow X$  which satisfies  $C(\gamma) = C(\alpha) * C(\beta)$  at each crossing, where  $\alpha, \gamma \in \Sigma$  are under-arcs on the right and left of the over-arc  $\beta \in \Sigma$ , respectively. If an  $X$ -coloring uses only one color we say that it is *trivial*.

For example, the coloring by the dihedral quandle  $R_m$  is coincident with Fox's  $m$ -coloring. An Alexander quandle coloring  $C$  satisfies  $C(\gamma) = tC(\alpha) + (1 - t)C(\beta)$  and  $C(\alpha) = t^{-1}C(\gamma) + (1 - t^{-1})C(\beta)$  at each crossing, where  $\alpha, \beta$  and  $\gamma$  are above-mentioned.


Let  $D$  be a diagram of an oriented knot  $K$  with a prime determinant  $p$ ,  $\Delta_K(t)$  be the Alexander polynomial of  $K$ , and  $\Sigma = \{\alpha_i\}_i$  be the set of arcs of  $D$ . Since the determinant  $p$  is equal to  $|\Delta_K(-1)|$ , an  $R_p$ -coloring of  $D$  corresponds to a coloring by  $\Lambda/(\Delta_K(t), t + 1) = \mathbb{Z}/(\Delta_K(-1))$ . For any color  $f_i(t)$  of  $\alpha_i$  colored by  $\Lambda/(\Delta_K(t))$ , there is a color  $a_i$  of  $\alpha_i$  colored by  $R_p$ , such that  $f_i(-1) = a_i \pmod{p}$ . If  $a_i$  is not equal to  $a_j$  in  $\mathbb{Z}_p$  for  $i \neq j$ , then  $f_i(t)$  is not equal to  $f_j(t)$  in  $\Lambda/(\Delta_K(t))$ . Therefore, if the Kauffman-Harary conjecture is true then Conjecture 1.2, also, is true for any alternating knots with prime determinants. We remark that there is a knot with non-prime determinants whose Alexander polynomial is a prime element in the Laurent polynomial ring  $\mathbb{Z}[t, t^{-1}]$ . See Section 3.

### 3. ALEXANDER QUANDLE COLORINGS OF TWIST KNOTS.

In this section, we consider Alexander quandle colorings of twist knots.

The diagram  $D_n$  of an  $n$ -twist knot  $K_n$  is pictured in the left of Figure 1, where  $|n|$  is the number of crossings in the "twist" part. The twists are right-handed if  $n > 0$  and left-handed if  $n < 0$ . The left of Figure 1 shows the case  $n > 0$ . We orient  $K_n$  by the orientation indicated in the left of Figure 1.

Let  $\Lambda/J$  be an Alexander quandle. We color the arcs of the "twist" part in the diagram  $D_n$  by  $a_0, b_0, b_1, \dots, b_{|n|} \in \Lambda/J$  as shown in the right of Figure 1. By the definition of Alexander quandle colorings, the relations between these colors are

FIGURE 1. The diagram  $D_n$  of  $K_n$ 

described by

$$\begin{pmatrix} b_{i-1} \\ b_i \end{pmatrix} = \begin{cases} \begin{pmatrix} 0 & 1 \\ t & 1-t \end{pmatrix} \begin{pmatrix} b_{i-2} \\ b_{i-1} \end{pmatrix} & \text{if } i \text{ is even,} \\ \begin{pmatrix} 0 & 1 \\ t^{-1} & 1-t^{-1} \end{pmatrix} \begin{pmatrix} b_{i-2} \\ b_{i-1} \end{pmatrix} & \text{if } i \text{ is odd,} \end{cases}$$

without regard to the sign of  $n$ , where  $i = 0, 1, \dots, n$  and  $b_{-1} = a_0$ . By induction, we have

$$(1) \quad \begin{pmatrix} b_{i-1} \\ b_i \end{pmatrix} = \frac{1}{2} \begin{pmatrix} i(t^{-1} - 1) + 2 & i(1 - t^{-1}) \\ i(t^{-1} - 1) & i(1 - t^{-1}) + 2 \end{pmatrix} \begin{pmatrix} a_0 \\ b_0 \end{pmatrix}$$

if  $i$  is even, and

$$(2) \quad \begin{pmatrix} b_{i-1} \\ b_i \end{pmatrix} = \frac{1}{2} \begin{pmatrix} (i-1)(t^{-1} - 1) & (i-1)(1 - t^{-1}) + 2 \\ (i+1)(t^{-1} - 1) + 2 & (i+1)(1 - t^{-1}) \end{pmatrix} \begin{pmatrix} a_0 \\ b_0 \end{pmatrix}$$

if  $i$  is odd. Furthermore, the colorings  $a_0, b_0, b_{|n|-1}, b_{|n|}$  of the four arcs in the “clasp” part of  $D_n$  have the following relations:

$$(3) \quad b_0 = \begin{cases} t^{-1}a_0 + (1 - t^{-1})b_n & \text{if } n \text{ is positive, even,} \\ ta_0 + (1 - t)b_n & \text{if } n \text{ is positive, odd,} \\ ta_0 + (1 - t)b_{|n|-1} & \text{if } n \text{ is negative, even,} \\ t^{-1}a_0 + (1 - t^{-1})b_{|n|-1} & \text{if } n \text{ is negative, odd,} \end{cases}$$

and

$$(4) \quad b_{|n|-1} = \begin{cases} t^{-1}b_n + (1 - t^{-1})a_0 & \text{if } n \text{ is positive,} \\ tb_{|n|} + (1 - t)b_0 & \text{if } n \text{ is negative.} \end{cases}$$

**Lemma 3.1.** *Assume that the ring  $\Lambda/J$  is an integral domain. The diagram  $D_n$  admits a non-trivial  $\Lambda/J$ -coloring if and only if it holds that*

$$\begin{cases} n(t^{-1} - 2 + t) = 2 & \text{if } n \text{ is positive, even,} \\ t^{-1} + \frac{1}{2}(n+1)(t^{-2} - 2t^{-1} + 1) = 0 & \text{if } n \text{ is positive, odd,} \\ |n|(t^{-1} - 2 + t) = -2 & \text{if } n \text{ is negative, even,} \\ t^{-1} - \frac{1}{2}(|n|-1)(t^{-2} - 2t^{-1} + 1) = 0 & \text{if } n \text{ is negative, odd.} \end{cases}$$

*Proof.* We assume that  $n$  is positive, even. From the relations (1),(3) and (4), we obtain  $(a_0 - b_0)(n(t^{-1} - 2 + t) - 2) = 0$ . If the color  $a_0$  is equal to the color  $b_0$  then  $D_n$  has nothing but trivial  $\Lambda/J$ -colorings. Since  $\Lambda/J$  is an integral domain,  $D_n$  admits a non-trivial  $\Lambda/J$ -coloring if and only if it holds that  $n(t^{-1} - 2 + t) = 2$ .

In the same way, we can prove this lemma for other cases.  $\square$

The Alexander polynomial  $\Delta_{K_n}(t)$  of the twist knot  $K_n$  is equal to

$$(5) \quad \begin{cases} \frac{n}{2}(1 - 2t + t^2) - t & \text{if } n \text{ is positive, even,} \\ t + \frac{1}{2}(n + 1)(1 - 2t + t^2) & \text{if } n \text{ is positive, odd,} \\ \frac{|n|}{2}(1 - 2t + t^2) + t & \text{if } n \text{ is negative, even,} \\ t - \frac{1}{2}(|n| - 1)(1 - 2t + t^2) & \text{if } n \text{ is negative, odd,} \end{cases}$$

up to multiplication by a unit  $\pm t^{\pm k}$ . There is an integer  $n$  such that, although the determinant  $|\Delta_{K_n}(-1)|$  is not prime, the Alexander polynomial  $\Delta_{K_n}(t)$  is prime in the Laurent polynomial ring  $\Lambda = \mathbb{Z}[t, t^{-1}]$ , that is, the ring  $\Lambda/(\Delta_{K_n}(t))$  is an integral domain. For example, if  $n$  is odd then  $\Delta_{K_n}(t)$  is always prime. We have the following proposition from Lemma 3.1 and (5).

**Proposition 3.2.** *For any integer  $n$  the diagram  $D_n$  of the twist knot  $K_n$  admits a non-trivial  $\Lambda/(\Delta_{K_n}(t))$ -coloring.*

We consider the case that  $n$  is positive, that is, we suppose that the diagram  $D_n$  is alternating. Assume that  $D_n$  is colored by a non-trivial  $\Lambda/(\Delta_{K_n}(t))$ -coloring. If the color  $b_l$  is equal to the color  $b_m$  for integers  $l, m$  ( $-1 \leq l, m \leq n$ ) then from the relations (1), (2) we obtain  $l = m$ . In other words, different arcs of  $D_n$  are colored by different colors. Accordingly, we have the following theorem.

**Theorem 3.3.** *Conjecture 1.2 is true for twist knots.*

In the same way, we can prove that for a negative integer  $n$ , that is, for a non-alternating diagram  $D_n$ , every non-trivial  $\Lambda/(\Delta_{K_n}(t))$ -coloring of  $D_n$  assigns different colors to different arcs of  $D_n$ . Possibly we may remove the condition that “a diagram is alternating” from Conjecture 1.2. At the present the author does not know a counterexample.

## REFERENCES

- [1] M. Asaeda, J. Przytycki, and A. Sikora, *Kauffman-Harary conjecture holds for Montesinos knots*, to appear in *J. Knot Theory Ramifications*.
- [2] R. H. Fox, *A quick trip through knot theory*, *Topology of 3-manifolds and related topics* (Georgia, 1961), Prentice-Hall, 120–167.
- [3] F. Harary, L. H. Kauffman, *Knots and Graphs I - Arc Graphs and Colorings*, *Advances in Applied Math.* **22** (1999), 312–337.
- [4] D. Joyce, *A classifying invariant of knots, the knot quandle*, *J. Pure Appl. Alg.* **23** (1982), 37–65.
- [5] S. Matveev, *Distributive groupoids in knot theory* (Russian), *Math. USSR-Sbornik* **23** (1982), 78–88.

DIVISION OF MATHEMATICAL SCIENCES AND PHYSICS, GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, CHIBA UNIVERSITY, 1-33 YAYOI-CHO, INAGE-KU, CHIBA, 263-8522, JAPAN

*E-mail address:* xasami@g.math.s.chiba-u.ac.jp